

# 2020

## Disease Risk Spray Schedules

Field Name \_\_\_\_\_

Planting Date \_\_\_\_\_

| PROGRAMS         | LEAF SPOT                 |                           | LEAF SPOT / WHITE MOLD / LIMB ROT          |  |  |   | LEAF SPOT                 |
|------------------|---------------------------|---------------------------|--|--|--|---|---------------------------|
|                  | 30                        | 45                        | 60   | 75   | 90   | 105   | 120                       |
| DAP <sup>1</sup> |                           |                           |  |  |  |   |                           |
| LOW RISK         | Chlorothalonil<br>1.5 pts | Chlorothalonil<br>1.5 pts | UMBRA 26 fl oz +<br>Chlorothalonil 1 pt    | Chlorothalonil<br>1.5 pts  | UMBRA 26 fl oz +<br>Chlorothalonil 1 pt    | Chlorothalonil<br>1.5 pts                       | Chlorothalonil<br>1.5 pts |
| MEDIUM RISK      | Chlorothalonil<br>1.5 pts | Chlorothalonil<br>1.5 pts | UMBRA 14-18 fl oz +<br>Chlorothalonil 1 pt | UMBRA 14-18 fl oz +<br>Chlorothalonil 1 pt                                 | UMBRA 14-18 fl oz +<br>Chlorothalonil 1 pt | UMBRA 14-18 fl oz +<br>Chlorothalonil 1 pt      | Chlorothalonil<br>1.5 pts |
| HIGH RISK        | Priaxor 6 fl oz           |                           | UMBRA 36 fl oz +<br>Chlorothalonil 1 pt    | Tebuconazole 7.2 fl oz +<br>Chlorothalonil 1 pt<br>OR<br>Priaxor 6-8 fl oz | UMBRA 36 fl oz +<br>Chlorothalonil 1 pt    | Tebuconazole 7.2 fl oz +<br>Chlorothalonil 1 pt | Chlorothalonil<br>1.5 pts |


<sup>1</sup>Days After Planting.

Notes: • Use higher rate of UMBRA if white mold risk increases to High Risk category.

- UMBRA controls soil-borne diseases (*Sclerotium rolfsii* – white mold/Southern blight; *Rhizoctonia solani* – limb rot) and foliar diseases (early and late leaf spot; peanut rust; web blotch).
- One pint of chlorothalonil should be used with all applications of UMBRA to reduce risk of resistance and to enhance leaf spot control.

See reverse side to assess the Peanut Disease Risk Index developed by:

UNIVERSITY OF GEORGIA      UNIVERSITY OF FLORIDA      AUBURN UNIVERSITY      MISSISSIPPI STATE UNIVERSITY      CLEMSON UNIVERSITY



**NICHINO**  
AMERICA